

OMIM
Online Mendelian Inheritance in Man

Johns Hopkins University

PubMed Nucleotide Protein Genome Structure PMC Taxonomy OMIM

Search for

Limits Preview/Index History Clipboard Details

Show:

***600355**

GeneTests, Links

BACULOVIRAL IAP REPEAT-CONTAINING PROTEIN 1; BIRC1**Alternative titles; symbols****NEURONAL APOPTOSIS INHIBITORY PROTEIN; NAIP**Gene map locus [5q12.2-q13.3](#)**TEXT**

In a search for the gene causing spinal muscular atrophy (SMA; [253300](#)), [Roy et al. \(1995\)](#) isolated a gene on chromosome 5q13.1, of which the first 2 coding exons were deleted in approximately 67% of type I SMA chromosomes compared with 2% of non-SMA chromosomes. One model of SMA pathogenesis invokes an inappropriate persistence of motor neuron apoptosis, which is a normally occurring phenomenon in development. Consistent with this hypothesis, the novel gene was labeled 'neuronal apoptosis inhibitory protein' (NAIP) and its function was supported by the finding that it contains domains with sequence similarity to IAPs, baculovirus proteins that inhibit virally induced insect cell apoptosis. The presence of a variable number of copies of truncated and internally deleted versions of the NAIP gene (pseudogenes) was thought to be a possible factor in the genesis of SMA. This situation was compared to that of the genes on 6p that code for steroid 21-hydroxylase (CYP21) deficiency ([201910](#)); through the processes of unequal crossing-over or gene conversion, mutations in the functional CYP21 gene can result. [Roy et al. \(1995\)](#) raised the possibility that NAIP functions in concert with SMN ([600354](#)) mutations in causing spinal muscular atrophy. 💡

Although NAIP deletions are more frequently observed in patients affected by the acute form of SMA, it is not possible to establish an unambiguous correlation between deletion size and clinical severity. [Novelli et al. \(1997\)](#) investigated the effects of gender on the association between NAIP gene deletion and disease severity. No significant relationship between deletion size and clinical phenotype was observed among male patients, whereas in females the absence of NAIP was strongly associated with a severe phenotype (p less than 0.0001). SMA I was found in 75.6% of females and only 52.5% of males lacking NAIP. These results provided a possible molecular explanation for the sex-dependent phenotypic variation observed in SMA patients. 💡

[DiDonato et al. \(1997\)](#) mapped the mouse homolog of NAIP to chromosome 13 in a region

showing conserved synteny with human 5q13.

Liston et al. (1996) demonstrated that expression of NAIP in mammalian cells inhibits apoptosis induced by a variety of signals.

NAIP, HIAP1 (601721), HIAP2 (601712), XIAP (300079), BIRC5 (603352), and BIRC6 (605638) are members of the mammalian inhibitors of apoptosis family and contain an N-terminal domain with 1 to 3 imperfect repeats of an approximately 65-amino acid domain named the baculovirus IAP repeat (BIR) motif. Gotz et al. (2000) identified 6 mouse Naip genes which were expressed in a broad range of tissues. Using a neurite outgrowth assay in rat pheochromocytoma PC12 cells, they observed that Naip overexpression impaired nerve growth factor (NGF)-induced neurite outgrowth. The BIR motifs of Naip (residues 1-345) were not required for this effect. However, the BIR domains of Naip were essential to prevent apoptosis in PC12 cells after NGF deprivation or tumor necrosis factor- α receptor (TNFAR; 191190) stimulation. Expression of full-length but not BIR-deleted Naip protected against cell death. This correlated with reduced activity of the cell death effector protease, caspase-3 (600636), in lysates of Naip-PC12 cells. The authors hypothesized that dysregulation of cellular differentiation and/or caspase suppression may contribute to motoneuron dysfunction and cell death in spinal muscular atrophy where NAIP is mutated. 🧠

ANIMAL MODEL

In inbred mouse strains, permissiveness to intracellular replication of *Legionella pneumophila* is controlled by a single locus (Lgn1), which maps to a region within distal chromosome 13 that contains multiple copies of the Birc1 gene. Genomic BAC clones from the critical interval were transferred into transgenic mice to complement functionally the Lgn1-associated susceptibility of A/J mice to *L. pneumophila*. Diez et al. (2003) found that 2 independent BAC clones that rescued susceptibility had an overlapping region of 56 kb in which the entire Lgn1 transcript must lie. The full-length transcript of Birc1e (also called Naip5) is coded in this region. The results indicated a role for Birc1e in macrophage resistance to *L. pneumophila* infection. BIRC1 proteins are members of the inhibitor of apoptosis protein (IAP) family, structurally defined by baculovirus inhibitor of apoptosis repeat (BIR) domains implicated in protein-protein interactions. An antiapoptotic effect of BIRC1 has been described. Induction of apoptosis seems to be important for pathogenesis of *L. pneumophila* in human macrophages in vitro. 🧠

REFERENCES

1. DiDonato, C. J.; Nadeau, J. H.; Simard, L. R. :
The mouse neuronal apoptosis inhibitory protein gene maps to a conserved syntenic region of mouse chromosome 13. *Mammalian Genome* 8: 222 only, 1997.
PubMed ID : [9069126](#)
2. Diez, E.; Lee, S.-H.; Gauthier, S.; Yaraghi, Z.; Tremblay, M.; Vidal, S.; Gros, P. :
Birc1e is the gene within the Lgn1 locus associated with resistance to *Legionella pneumophila*. *Nature Genet.* 33: 55-60, 2003.
PubMed ID : [12483212](#)

3. Gotz, R.; Karch, C.; Digby, M. R.; Troppmair, J.; Rapp, U. R.; Sendtner, M. :
The neuronal apoptosis inhibitory protein suppresses neuronal differentiation and apoptosis in PC12 cells. *Hum. Molec. Genet.* 9: 2479-2489, 2000.
PubMed ID : [11030753](#)
4. Liston, P.; Roy, N.; Tamai, K.; Lefebvre, C.; Baird, S.; Cherton-Horvat, G.; Farahani, R.; McLean, M.; Ikeda, J.-E.; MacKenzie, A.; Korneluk, R. G. :
Suppression of apoptosis in mammalian cells by NAIP and a related family of IAP genes. *Nature* 379: 349-353, 1996.
PubMed ID : [8552191](#)
5. Novelli, G.; Semprini, S.; Capon, F.; Dallapiccola, B. :
A possible role of NAIP gene deletions in sex-related spinal muscular atrophy phenotype variation. *Neurogenetics* 1: 29-30, 1997.
PubMed ID : [10735271](#)
6. Roy, N.; Mahadevan, M. S.; McLean, M.; Shutler, G.; Yaraghi, Z.; Farahani, R.; Baird, S.; Besner-Johnston, A.; Lefebvre, C.; Kang, X.; Salih, M.; Aubry, H.; Tamai, K.; Guan, X.; Ioannou, P.; Crawford, T. O.; de Jong, P. J.; Surh, L.; Ikeda, J.-E.; Korneluk, R. G.; MacKenzie, A. :
The gene for neuronal apoptosis inhibitory protein is partially deleted in individuals with spinal muscular atrophy. *Cell* 80: 167-178, 1995.
PubMed ID : [7813013](#)

CONTRIBUTORS

Victor A. McKusick – updated : 12/18/2002
George E. Tiller – updated : 1/16/2001
Rebekah S. Rasooly – updated : 2/22/1999
Victor A. McKusick – updated : 9/12/1997
Victor A. McKusick – updated : 4/15/1997

CREATION DATE

Victor A. McKusick : 1/27/1995

EDIT HISTORY

alopez : 12/18/2002
terry : 12/18/2002
carol : 2/12/2001
carol : 1/26/2001
mcapotos : 1/26/2001
mcapotos : 1/19/2001
mcapotos : 1/16/2001
mcapotos : 1/16/2001
mgross : 9/15/2000
alopez : 2/22/1999
alopez : 2/22/1999

terry : 9/12/1997
jenny : 4/15/1997
terry : 4/10/1997
mark : 10/8/1996
mark : 7/22/1996
mimadm : 9/23/1995
terry : 1/27/1995

Copyright © 1966-2003 Johns Hopkins University

Display Detailed Show: 20 Send to Text

[Disclaimer](#) | [Write to the Help Desk](#) | [Privacy Policy](#)
[NCBI](#) | [NLM](#) | [NIH](#)

Search notes for 09830338
12/19/2003

SYSTEM:OS - DIALOG OneSearch

File 34:SciSearch(R) Cited Ref Sci 1990-2003/Dec W2

(c) 2003 Inst for Sci Info

File 35:Dissertation Abs Online 1861-2003/Nov

(c) 2003 ProQuest Info&Learning

File 399:CA SEARCH(R) 1967-2003/UD=13925

(c) 2003 American Chemical Society

*File 399: Use is subject to the terms of your user/customer agreement.

Alert feature enhanced for multiple files, etc. See HELP ALERT.

Set Items Description

--- -----

?s naip and monoclonal

309 NAIP

214338 MONOCLONAL

S1 4 NAIP AND MONOCLONAL

?rd

...completed examining records

S2 4 RD (unique items)

File 5:Biosis Previews(R) 1969-2003/Dec W2

(c) 2003 BIOSIS

Set Items Description

S1 201 NAIP

S2 0 S1 AND MONOCLON?

09/830, 338
Search notes 12-19-00Product Number: **N6648**Product Name: **Anti-NAIP antibody produced in rabbit**[Register or Login
for Pricing
Click Here](#)[Product Information](#)

Synonyms: MDL number: MFCD01322381

[Description](#)

Storage Temp: -20°C

[Certificate of Analysis](#)[MSDS](#)[Datasheet](#)[Structure Image](#)[Questions](#)**Comments:**

Affinity isolated antibody, Lyophilized powder

Species reactivity: human

[Print Preview](#)[Bulk Quote](#)[Ask A Scientist](#)**Immunogen:** synthetic peptide corresponding to amino acids 473-490 of human NAIP (neuronal apoptosis inhibitory protein) sequence, conjugated to KLH.**Physical form:** Lyophilized from a 0.2 µm filtered solution in phosphate buffered saline**Extended specifications****Application(s)**

Immunoblotting

10 µg/mL

Components:**Related Product(s):**[help](#) | [privacy](#) | [technical library](#) | [search](#) | [home](#)
[terms and conditions](#) | [contract manufacturing](#)

© 2003 Sigma-Aldrich Co. Reproduction forbidden without permission.

Sigma-Aldrich brand products are sold exclusively through Sigma-Aldrich, Inc. Best viewed in IE5 or higher

This is the html version of the file <http://bst.portlandpress.com/bst/029/0296740025.pdf>.

Google automatically generates html versions of documents as we crawl the web.

To link to or bookmark this page, use the following url: [http://www.google.com/search?](http://www.google.com/search?q=cache:JNeV9oZtwzUJ:bst.portlandpress.com/bst/029/0296740025.pdf+monoclonal+naip&hl=en&ie=UTF-8)

[q=cache:JNeV9oZtwzUJ:bst.portlandpress.com/bst/029/0296740025.pdf+monoclonal+naip&hl=en&ie=UTF-8](http://www.google.com/search?q=cache:JNeV9oZtwzUJ:bst.portlandpress.com/bst/029/0296740025.pdf+monoclonal+naip&hl=en&ie=UTF-8)

Google is not affiliated with the authors of this page nor responsible for its content.

These search terms have been highlighted: **monoclonal naip**

Page 1

25 The neuronal apoptosis inhibitory protein (NAIP) is a membrane-associated cytoplasmic protein in rabbit brain.

T.A. Lynch and G.E. Morris

MRIC Biochemistry Group, North East Wales Institute, Wrexham LL11 2AW, UK.

The Neuronal Apoptosis Inhibitory Protein (NAIP) gene was originally identified as a candidate gene for Spinal Muscular Atrophy (SMA). This disease is characterized by motor neuron degeneration, with evidence for an apoptotic mechanism. NAIP shares significant homology with two baculoviral inhibitor of apoptosis proteins, Cp-IAP and Op-IAP.

We have produced a panel of three monoclonal antibodies against a region of NAIP containing the baculovirus inhibitor of apoptosis repeats (BIRs). The antibodies recognized a protein of the predicted size (150kDa) in all brain regions tested and at lower levels in spinal cord and peripheral nerve. NAIP was undetectable in liver, heart and skeletal muscle. Immunohistochemistry of rabbit brain showed NAIP expression in the cytoplasm of neuronal cells, especially Purkinje neurons of the cerebellum and their dendritic processes. Subcellular fractionation of rabbit brain showed that the 150kDa NAIP protein sediments in heavy and light membrane fractions and is extractable with non-ionic detergent. The results suggest that NAIP is associated with internal membranes of the endoplasmic reticulum via a transmembrane sequence (aa 479-496). The N-terminal region, containing BIR domains and nucleotide-binding site, projects into the cytoplasm.

Supported by the Muscular Dystrophy Association (USA).

25 The neuronal apoptosis inhibitory protein (NAIP) is a membrane-associated cytoplasmic protein in rabbit brain.

T.A. Lynch and G.E. Morris

MRIC Biochemistry Group, North East Wales Institute, Wrexham LL11 2AW, UK.

The Neuronal Apoptosis Inhibitory Protein (NAIP) gene was originally identified as a candidate gene for Spinal Muscular Atrophy (SMA). This disease is characterized by motor neuron degeneration, with evidence for an apoptotic mechanism. NAIP shares significant homology with two baculoviral inhibitor of apoptosis proteins, Cp-IAP and Op-IAP.

We have produced a panel of three monoclonal antibodies against a region of NAIP containing the baculovirus inhibitor of apoptosis repeats (BIRs). The antibodies recognized a protein of the predicted size (150kDa) in all brain regions tested and at lower levels in spinal cord and peripheral nerve. NAIP was undetectable in liver, heart and skeletal muscle. Immunohistochemistry of rabbit brain showed NAIP expression in the cytoplasm of neuronal cells, especially Purkinje neurons of the cerebellum and their dendritic processes. Subcellular fractionation of rabbit brain showed that the 150kDa NAIP protein sediments in heavy and light membrane fractions and is extractable with non-ionic detergent. The results suggest that NAIP is associated with internal membranes of the endoplasmic reticulum via a transmembrane sequence (aa 479-496). The N-terminal region, containing BIR domains and nucleotide-binding site, projects into the cytoplasm.

Supported by the Muscular Dystrophy Association (USA).



Send note
SIGMA

09/830, 338

ANTI-NAIP

Developed in Rabbit
Affinity Isolated Antibody

Product Number **N 6648**

Product Description

Anti-NAIP is developed in rabbit using a synthetic peptide conjugated to KLH as immunogen. This peptide corresponds to amino acids 473-490 (YLSLSSTRPDE-GLASLIC) of human NAIP. This antibody is affinity-purified human NAIP using peptide affinity chromatography.

Anti-NAIP detects human NAIP in extracts from *Sf 21* insect cells expressing recombinant human NAIP by immunoblotting.

NAIP (neuronal apoptosis inhibitor protein) is a member of the IAPs (inhibitor of apoptosis proteins) that function in cell death pathways to inhibit programmed cell death.¹ Human NAIP has a calculated molecular mass of approximately 140 kDa. The IAPs share one to three copies of an approximately 70 amino acid sequence motif, BIR (baculovirus IAP repeat).² These BIR regions promote protein-protein interactions with caspases as well as with members of the TRAF family of signal molecules.³

The first human IAP identified was NAIP, discovered based on its association with a neurodegenerative disorder. Mutations and deletions in the NAIP gene locus is a contributing factor in spinal muscular atrophy.⁴ Co-expression of NAIP and hippocalcin protects neurons against calcium-induced cell death in caspase-3-activated and non-activated pathways.⁵ NAIP is strongly expressed in anterior horn and motor cortex neurons of the normal brain. It is also found in human fetal neurons and in adult choroid plexus cells.⁶

Reagent

Anti-XIAP is supplied as 100 µg of antibody lyophilized from a 0.2 µm filtered solution in phosphate buffered saline.

Preparation Instructions

To one vial of lyophilized powder, add 0.1 ml of 0.2 µm-filtered solution of phosphate-buffered saline (PBS) containing 0.02% sodium azide to produce a 1.0 µg/ml stock solution of antibody.

Product Information

Storage/Stability

Prior to reconstitution, store at -20° C. The reconstituted product may be stored at 2-8° C for at least one month. For prolonged storage, freeze in working aliquots at -20° C. Avoid repeated freezing and thawing.

Product Profile

The recommended working concentration is 1.0 µg/ml for immunoblotting using extracts from *Sf 21* cells expressing human NAIP by chemiluminescent detection. The ability of the antibody to blot endogenous NAIP in cell extracts is not known.

Note: In order to obtain best results in different techniques and preparations we recommend determining optimal working dilutions by titration.

References

1. Wagenknecht, B., et al., Expression and biological activity of X-linked inhibitor of apoptosis (XIAP) in human malignant glioma. *Cell Death Differ.*, **6**, 370-376 (1999).
2. Sun, C., et al., NMR structure and mutagenesis of the inhibitor-of-apoptosis protein XIAP. *Nature*, **401**, 818-822 (1999).
3. Roy, N., et al., The c-IAP-1 and c-IAP-2 proteins are direct inhibitors of specific caspases. *EMBO J.*, **16**, 6914-6925 (1997).
4. Roy, N., et al., The gene for neuronal apoptosis inhibitory protein is partially deleted in individuals with spinal muscular atrophy. *Cell*, **80**, 167-178 (1995).
5. Mercer, E.A. et al., NAIP interacts with hippocalcin and protects neurons against calcium-induced cell death through caspase-3-dependent and-independent pathways. *EMBO J.*, **19**, 3597-3607 (2000).
6. Pari, G., et al., Immunolocalization of NAIP in the human brain and spinal cord. *Neuroreport*, **11**, 9-14 (2000).

KAA 10/00

Sigma brand products are sold through Sigma-Aldrich, Inc.

Sigma-Aldrich, Inc. warrants that its products conform to the information contained in this and other Sigma-Aldrich publications. Purchaser must determine the suitability of the product(s) for their particular use. Additional terms and conditions may apply. Please see reverse side of the invoice or packing slip.